

Prostomidae (Coleoptera: Tenebrionoidea) – a New Family to the Fauna of Bulgaria

*Borislav V. Guéorguiev*¹

¹ National Museum of Natural History – Sofia, 1 Blvd. Tzar Osvoboditel, 1000 Sofia, Bulgaria, E-mail: bobivg@yahoo.com

Abstract: Family Prostomidae, genus *Prostomis*, and species *P. mandibularis* are recorded for first time from Bulgaria. *P. mandibularis*, characteristic of decaying heartwood of rotting wood lying undisturbed on forest floors, is a species of conservation importance and indicator of use for assessment of natural old and well-preserved broad-leaved woodlands.

Key words: Coleoptera, *Prostomis mandibularis*, Bulgaria, first record.

Introduction

The number of families from order Coleoptera which occur in Bulgaria is well-known. Following the classification of FAUNA EUROPAEA WEB SERVICE 2004 (ALONSO-ZARAZAGA 2004, AUDISIO 2004)¹, we know data for representatives from 117 families there; this account includes only taxa native to the fauna of Europe. Up to now, we had no records only for species from Rhipiceridae (Dascilloidea), Endecatomidae (Bostrichoidea), Phloiophilidae (Cleroidea), Phloeostichidae (Cucujoidea), Prostomidae, Boridae (Tenebrionoidea), and Raymondionymidae (Curculionoidea) (GUÉORGUIEV *et al.* 2008)².

Recent pitfall collecting in old, predominantly *Castanea sativa* Miller forests revealed a further new family to the fauna of Bulgaria. The purpose of the present report is to announce this finding.

Prostomidae C.G. THOMSON, 1859

***Prostomis* LATREILLE, 1825**

***Prostomis mandibularis* FABRICIUS, 1801**

(Fig. 1)

Material studied

Belasitsa Mountain: N41.371754 E23.185813, below Belasitsa Chalet, 700 m, 14.IX-30.X.2009, 1 specimen, pitfall traps, leg. B. Guéorguiev & P. Mitov; N41.352124 E23.204887, 2.7 km SE Belasitsa Chalet, 800 m, 26.III-18.V.2010, 1 specimen / 18.V-4.VII.2010, 1 specimen, pitfall traps, leg. B. Guéorguiev & Ch. Deltshv; N41.363556 E23.210198, ca. 4 km S of Petrich Town, 800 m, 29.III-19.V.2010, 1 specimen, pitfall traps, leg. B. Guéorguiev & Ch. Deltshv; N41.354371 E23.204756, ca. 4 km S of Petrich Town, 750 m, 18.V-4.VII.2010, 1 specimen, leg. B. Guéorguiev &

¹ The discrepancy in the family number in the present report and GUÉORGUIEV *et al.* (2008) is due to the use of differing classifications.

² Cerophytidae has been also included among the family group-taxa unknown to Bulgaria, but in fact it was noted for the country the year before (BOČÁK 2007).



Fig. 1. Habitus of *Prostomis mandibularis* Fabricius, Belasitsa Mt., Bulgaria. Scale 1 mm.

Ch. Deltshev. The material is preserved in the collection of the National Museum of Natural History, Sofia. Data for the forest communities and tree age are shown in Table 1.

Faunistic notes and biology

The family Prostomidae (jugular-horned beetles) comprises 28 species arranged in one extinct (*Vetuprostomis* ENGEL & GRIMALDI, 2008) and two extant genera (*Prostomis* LATREILLE, 1825 and *Dryocora* PASCOE, 1868) (SCHAWALLER 2003, PARK, AHN 2005, ENGEL, GRIMALDI 2008). The species from the extant genera are distributed worldwide,

excluding only the Neotropic Region and most part of Africa (SCHAWALLER 2003). Adult members of this group are readily recognized by combination of the following characters (Fig. 1): body elongated, parallel-sided and strongly flattened; colour red to reddish-brown; prognathic head having large, robust and strongly projecting mandibles (dorsal view) as well as large and projecting forward jugular processes (ventral view); antennae relatively short, not exceeding posterior margin of pronotum with last three articles slightly clubbed; pronotum as wide as head; elytra as wide as pronotum; tarsal formulae 4-4-4.

In this work, we report *P. mandibularis* for first time for the fauna of Bulgaria; the species range includes Europe, the Caucasus, North Turkey, and North Iran (SCHAWALLER 2003). This curiously flattened (in both stages larva and imago) beetle is xylophagous and characteristic of decaying heartwood of rotting wood lying undisturbed on forest floors. Like other members of the family (GROVE 2007), the species is supposedly feeding ‘mudguts’³ and tunnelling within and between the ‘mudguts’ and surrounding more solid wood.

P. mandibularis is presumably entirely dependent upon veteran trees⁴ as it inhabits decaying heartwood, a very specific habitat type which is already highly fragmented and subject to continuing significant decline in Europe. According to the recent European Red List of Saproxylic Beetle, it is listed as ‘Near Threatened’ (NIETO, ALEXANDER 2010). Therefore, we have an indicator species on European scale of use for assessment of natural old and well-preserved broad-leaved woodland communities.

Acknowledgements: The material has been collected as results of N BG 0031 – EEA FM ‘State and prospects of the *Castanea sativa* population in Belasitsa mountain: climate change adaptation, maintenance of biodiversity and sustainable ecosystem management’, a grant of the European Economic Area (EEA) и Norway Grants, and N BM06/07 ‘Assessment of forest communities of conservation importance based on invertebrate indicator groups in Belasitsa Mt. (Bulgaria and Republic of Macedonia),

³ According to some rotten wood type classification, the ‘mudguts’ is advanced stage of decay of the inner heartwood (YEE *et al.* 2001, YEE *et al.* 2006, GROVE 2007).

⁴ In Tasmania, another species from *Prostomis* was found only in mature trees more than 150 years old (HOPKINS *et al.* 2005).

Table 1. Localities and data for forest community and tree age.

Locality	Predominant tree species (tree age)
N41.371754 E23.185813	One generation <i>C. sativa</i> (100-150 years old); second generation of predominant <i>Quercus petraea</i> Liebl. (50-60 years old)
N41.352124 E23.204887	Predominant <i>C. sativa</i> (60-140 years old)
N41.363556 E23.210198	One generation <i>C. sativa</i> (100-150 years old); second generation of predominant <i>Fagus sylvatica</i> L. (50-60 years old)
N41.354371 E23.204756	Mixed forest <i>C. sativa</i> , <i>F. sylvatica</i> and <i>Quercus</i> spp. (60-140 years old)

a grant of the National Science Fund of the Bulgarian Ministry of Education and Science. Thanks due to Rostislav Bekchiev (National Museum of Natural

History – Sofia) for preparing Fig. 1 and Dr Tsvetan Zlatanov (Forestry Research Institute – Sofia) for providing data for forest community and tree age.

References:

- ALONSO-ZARAZAGA M. A. (Ed.) 2004. Coleoptera 1. Fauna Europea version 1.1, <http://www.faunaeur.org>
- AUDISIO P. (Ed.) 2004. Coleoptera 2. Fauna Europea version 1.1, <http://www.faunaeur.org>
- BOCÁK L. 2007. Cerophytidae. – In: Löbl I., A. Smetana (Eds.): Catalogue of Palaearctic Coleoptera. Vol. 4. Stenstrup: Apollo Books, 81.
- ENGEL M. S., D. A. GRIMALDI 2008. A jugular-horned beetle in Cretaceous amber from Myanmar (Coleoptera: Prostomidae). – *Alavesia*, **2**: 215-218.
- FAUNA EUROPAEA WEB SERVICE 2004. Fauna Europaea version 1.1, Available online at <http://www.faunaeur.org>
- GROVE S. J. 2007. Mudguts. – *The Tasmanian Naturalist*, **129**: 2-7.
- GUÉORGUIEV B., D. DOYCHEV and D. OVCHAROV 2008. Cucujidae (Coleoptera: Cucujoidea) – a new family to the fauna of Bulgaria. – *Historia naturalis bulgarica*, **19**: 93-97.
- HOPKINS A. J. M., K. S. HARRISON I, S. J. GROVE, T. J. WARDLAW and C. L. MOHAMMED 2005. Wood-decay fungi and saproxylic beetles associated with living *Eucalyptus obliqua* trees: early results from studies at the Warra LTER Site, Tasmania. – *Tasforests*, **16**: 111-126.
- NIETO A., K. N. A. ALEXANDER 2010. European Red List of Saproxylic Beetles. Luxembourg: Publications Office of the European Union.
- PARK J. S., K. J. AHN 2005. A taxonomic note on the little known family Prostomidae in Korea (Coleoptera: Tenebrionoidea). – *Entomological Research*, **35** (3): 169-171.
- SCHAWALLER W. 2003. New species and records of *Prostomis* Latreille, including the first fossil records from Baltic amber and a checklist of the species (Coleoptera: Prostomidae). – *Stuttgarter Beitrage zur Naturkunde, Serie A (Biologie)*, **650**: 1-11.
- YEE M., Z.-Q. YUAN and C. MOHAMMED 2001. Not just waste wood: decaying logs as key habitats in Tasmania's wet sclerophyll *Eucalyptus obliqua* production forests: the ecology of large and small logs compared. – *Tasforests*, **13**: 119-128.
- YEE M., S. J. GROVE, A. RICHARDSON and C. L. MOHAMMED. 2006. Brown rot in inner heartwood: why large logs support characteristic saproxylic beetle assemblages of conservation concern. – In: S. J. Grove & J. L. Hanula (Eds.). Insect biodiversity and dead wood. Proceedings of a symposium at the International Congress of Entomology, Brisbane, Australia, August 2004. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southern Research Station, Athens: 42-56.

Received: 10.09.2010
Accepted: 13.01.2011